

# MEDURAMAX

## COMPOSITION

Each kg contains: maduramycin ammonium  
10 g.

## DESCRIPTION

**MADURA-MAX** contains maduramycin, an ionophore antibiotic and anticoccidial. Maduramycin exerts its anticoccidial effect on the asexual forms of the protozoon (sporozoites, merozoites and falciform bodies). This interrupts the endogenous development of the parasite by interfering the ionic flux, especially potassium and sodium.

**MADURA-MAX** is effective against poultry coccidian infections even those coccidia resistant to other ionophore drugs.

## INDICATIONS

Prevention and treatment of coccidian infections in poultry (broiler, layers, replacement layers and breeders).

## CONTRAINDICATIONS

None described.

## ADVERSE EFFECTS

None described at the recommended dose.  
Overdose can cause decreased growth, leg weakness and even death in severe over dosage.

## SPECIAL PRECAUTIONS

Product intended to be mixed with feed. Do not use as is. Mix well with the feed to ensure a

homogeneous distribution. Avoid the contact with the skin and mucous. Use gloves, glasses and mask during the manipulation of the premix.

## ADMINISTRATION

By oral route, mixed with feed.  
500 g/1000 kg (5 ppm).

## TARGET SPECIES

Poultry (broilers, laying hens, replacement layers and breeders).

## WITHDRAWAL PERIOD

Poultry: meat, offal and eggs: five days.

## STORAGE CONDITIONS

Store below 30 °C.  
Store in a dry place, away from light.

## SIZES

25 Kg



OZONE

Rasquera-4, Tortosa-43500, (Tarragona) SPAIN

PH: (+34)977444975

[www.ozone.eu.com](http://www.ozone.eu.com)