

SALINO-MAX

COMPOSITION

Each kg contains: salinomycin 120 g.

DESCRIPTION

SALINO-MAX contains salinomycin, an ionophore antibiotic and anticoccidial. Salinomycin exerts its anticoccidial effect on the asexual forms of the oocyst (sporozoites, merozoites and falciform bodies). This interrupts the endogenous development of the parasite by interfering the ionic flux, especially potassium and sodium.

SALINO-MAX is effective against *Eimeria acervulina*, *Eimeria brunetti*, *Eimeria maxima*, *Eimeria mivati*, *Eimeria necatrix* and *Eimeria tenella*.

INDICATIONS

Prevention and treatment of coccidian infections in poultry (broiler, layers, replacement layers and breeders) and pigs.

CONTRAINDICATIONS

Not described.

ADVERSE EFFECTS

None described at the recommended dose. Do not use in target species off-label, especially horses.

SPECIAL PRECAUTIONS

Product intended to be mixed with feed. Do not use as is. Mix well with the feed to ensure a homogeneous distribution. Avoid the contact with the skin and mucous. Use gloves, glasses and mask during the manipulation of the premix.

ADMINISTRATION

By oral route, mixed with feed.

Poultry: 500 g/1000 kg feed.

Piglets less than 4 months of age: 250 – 500 g/1000 kg feed.

TARGET SPECIES

Poultry (broilers, and replacement pullets) and pigs (piglets).

WITHDRAWAL PERIOD

Meat, offal: zero days

Eggs: ten days.

STORAGE CONDITIONS

Store below 30 °C.

Store in a dry place, away from light.

SIZES

25 Kg



OZONE

Rasquera-4, Tortosa-43500, (Tarragona) SPAIN

PH: (+34)977444975

www.ozone.eu.com